European Journal of Histochemistry

SUPPLEMENTARY MATERIAL

DOI: <u>10.4081/ejh.2018.2866</u>

Diagnosis of sudden cardiac death due to early myocardial ischemia:

An ultrastructural and immunohistochemical study

Silvia Damiana Visonà,¹ Donatella Benati,² Maria Cristina Monti,³ Mirco Galiè,² Luisa Andrello,⁴ Andrea Frontini,⁵ Antonio Osculati¹

Correspondence: Silvia Damiana Visonà, Department of Public Health, Experimental and Forensic Medicine, via Forlanini 12, 27100 Pavia, Italy. Tel.: +39.320.6056485 - Fax: +39.0382.528052. E-mail: visona.silvia@gmail.com

Key words: Sudden death; early myocardial ischemia; ultrastructure; immunohistochemistry; Connexin 43.



¹Department of Public Health, Experimental and Forensic Medicine, Unit of Legal Medicine and Forensic Sciences, University of Pavia, Italy

²Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Italy

³Department of Public Health, Experimental and Forensic Medicine, Unit of Biostatistics and Clinical Epidemiology, University of Pavia, Italy

⁴Service of Legal Medicine, Canton Ticino, Bellinzona, Switzerland

⁵Department of Public Health, Experimental and Forensic Medicine, Unit of Normal Human Anatomy, University of Pavia, Italy

Supplementary Table 1. Case profiles (n=17). IVA= interventricular anterior branch of left coronary.

	Age	Sex	Coronary lesions	Main changes at traditional histology	Heart weight (g)	Resuscitation enacted	Post-mortem interval (hours)
1	51	M	IVA subocclusion	Areas of coagulative necrosis	450	Yes	44
2	70	F	Trivasal coronaric atherosclerosis	Areas of coagulative necrosis	530	Yes	195
3	62	M	IVA subocclusion	Areas of coagulative necrosis	450	No	21
4	39	M	No	"Wavy fibers", contraction bands	406	Yes	40
5	44	M	No (cocaine abuser)	"Wavy fibers", contraction bands	450	No	54
6	68	M	Trivasal coronaric atherosclerosis	Areas of coagulative necrosis	528	Yes	35
7	47	F	No	"Wavy fibers", contraction bands	290	Yes	11
8	82	M	Right coronaric subocclusion	Areas of coagulative necrosis	445	No	21
9	45	M	Subocclusion of common branch of left coronary artery	Areas of coagulative necrosis	465	No	27
10	51	F	Trivasal coronarosclerosis	Areas of coagulative necrosis	350	No	170



11	39	F	Subocclusion IVA	"Wavy fibers", contraction bands	320	Yes	80
12	74	M	Subocclusion of right cor + circumflex branch	Areas of coagulative necrosis	340	Yes	90
13	53	M	Thrombosis IVA, trivasal coronaric atherosclerosis	Areas of coagulative necrosis	440	No	30
14	63	M	No	"Wavy fibers", contraction bands, nuclear alterations	560	No	30
15	33	F	Yes (IVA 50%)	"Wavy fibers", contraction bands, nuclear alterations	370	Yes	36
16	78	M	Yes (right coronaric subocclusion)	Areas of coagulative necrosis	620	Yes	68
17	51	M	Yes (occlusions of IVA and Circumflex)	"Wavy fibers", contraction bands, nuclear alterations	600	No	96



Supplementary Table 2. Control profiles (n=10).

	Age	Sex	Cause of death	Heart weight (g)	Resuscitation enacted	Post-mortem interval (hours)
18	47	M	Gunshot wounds	390	No	42
19	60	F	Aortic dissection	300	Yes	135
20	56	M	Head and cervical trauma	330	No	32
21	41	M	Road traffic accident (decapitation)	300	No	16
22	43	M	Gunshot wounds	480	No	34
23	63	M	Fall from height	310	No	12
24	37	F	Fall from height	246	No	17
25	53	M	Road traffic accident	305	No	24
26	52	M	Gunshot wounds	350	No	56
27	39	F	Fall from height	250	No	22

