









11. Adarichev VA, Glant TT. Experimental spondyloarthropathies: animal models of ankylosing spondylitis. *Curr Rheumatol Rep* 2006;8:267-74.
12. Lories RJ. Animal models of spondyloarthritis. *Curr Opin Rheumatol* 2006;18: 342-6.
13. Braem K, Lories RJ. Insights into the pathophysiology of ankylosing spondylitis: contributions from animal models. *Joint Bone Spine* 2012;79:243-8.
14. Zhang Y, Shi S, Ciurli C, Poole AR. Animal models of ankylosing spondylitis. *Curr Rheumatol Rep* 2002; 4:507-12.
15. Oostveen J, Prevo R, den Boer J, van de Laar M. Early detection of sacroiliitis on magnetic resonance imaging and subsequent development of sacroiliitis on plain radiography. A prospective, longitudinal study. *J Rheumatol* 1999; 26:1953-8.
16. Blum U, Buitrago-Tellez C, Munding A, Krause T, Laubenberger J, Vaith P, et al. Magnetic resonance imaging (MRI) for detection of active sacroiliitis - a prospective study comparing conventional radiography, scintigraphy, and contrast enhanced MRI. *J Rheumatol* 1996;23: 2107-15.
17. Appel H, Loddenkemper C, Grozdanovic Z, Ebhardt H, Dreimann M, Hempfing A, et al. Correlation of histopathological findings and magnetic resonance imaging in the spine of patients with ankylosing spondylitis. *Arthritis Res Ther* 2006; 8:R143.
18. Ballet S, Conrath M, Fischer J, Kaneko T, Hamon M, Cesselin F. Expression and G-protein coupling of mu-opioid receptors in the spinal cord and dorsal root ganglia of polyarthritic rats. *Neuropeptides* 2003;37: 211-9.
19. Iwamoto T, Takasugi Y, Higashino H, Ito H, Koga Y, Nakao S. Antinociceptive action of carbamazepine on thermal hypersensitive pain at spinal level in a rat model of adjuvant-induced chronic inflammation. *J Anesth* 2011;25:78-86.
20. Gouret C, Mocquet G, Raynaud G. Use of Freund's adjuvant arthritis test in anti-inflammatory drug screening in the rat: value of animal selection and preparation at the breeding center. *Lab Anim Sci* 1976;26:281-7.
21. Dawson J, Gustard S, Beckmann N. High-resolution three-dimensional magnetic resonance imaging for the investigation of knee joint damage during the time course of antigen-induced arthritis in rabbits. *Arthritis Rheum* 1999;42:119-28.
22. Verdenius HH, Alma L. A quantitative study of decalcification methods in histology. *J Clin Pathol* 1958; 11:229-36.
23. Kalscheur VL. Bone and cartilage in osteoarthritis: a proteoglycan stain. *HistoLogic Techn Bull Histotechnol* 2001; XXXIV:14-5.
24. Gomori G. Aldehyde-fuchsin: a new stain for elastic tissue. *Am J Clin Pathol* 1950; 20:665-6.
25. Dijkstra CD, Dopp EA, van den Berg TK, Damoiseaux JG. Monoclonal antibodies against rat macrophages. *J Immunol Methods* 1994;174:21-3.
26. Damoiseaux JG, Dopp EA, Calame W, Chao D, MacPherson GG, Dijkstra CD. Rat macrophage lysosomal membrane antigen recognized by monoclonal antibody ED1. *Immunology* 1994;83:140-7.
27. Tan S, Yao J, Flynn JA, Yao L, Ward MM. Quantitative syndesmophyte measurement in ankylosing spondylitis using CT: longitudinal validity and sensitivity to change over 2 years. *Ann Rheum Dis* 2015;74:437-43.
28. Baeten D, Baraliakos X, Braun J, Sieper J, Emery P, van der Heijde D, et al. Anti-interleukin-17A monoclonal antibody secukinumab in treatment of ankylosing spondylitis: a randomised, double-blind, placebo-controlled trial. *Lancet* 2013;382: 1705-13.
29. Baraliakos X, Borah B, Braun J, Baeten D, Laurent D, Sieper J, et al. Long-term effects of secukinumab on MRI findings in relation to clinical efficacy in subjects with active ankylosing spondylitis: an observational study. *Ann Rheum Dis* 2016;75:408-12.

Non commercial use only