

Lu *et al.*²⁴ have found that the exogenous Wnt agonist BIO overcame the downregulation of SCs proliferation in *Bmi1*^{-/-} mice, providing a possibility of exogenous factors involved. There is more work waiting for us.

References

- Dayaratne MW, Vlajkovic SM, Lipski J, Thorne PR. Kolliker's organ and the development of spontaneous activity in the auditory system: implications for hearing dysfunction. *Biomed Res Int* 2014;2014:367939.
- Anniko M. Postnatal maturation of cochlear sensory hairs in the mouse. *Anat Embryol (Berl)* 1983;166:355-68.
- Tritsch NX, Yi E, Gale JE, Glowatzki E, Bergles DE. The origin of spontaneous activity in the developing auditory system. *Nature* 2007;450:50-5.
- Sendin G, Bourien J, Rassendren F, Puel JL, Nouvian R. Spatiotemporal pattern of action potential firing in developing inner hair cells of the mouse cochlea. *Proc Natl Acad Sci USA* 2014;111:1999-2004.
- He S, Yang J. Maturation of neurotransmission in the developing rat cochlea: immunohistochemical evidence from differential expression of synaptophysin and synaptobrevin 2. *Eur J Histochem* 2011;55:e2.
- Kennedy HJ. New developments in understanding the mechanisms and function of spontaneous electrical activity in the developing mammalian auditory system. *J Assoc Res Otolaryngol* 2012;13:437-45.
- Knipper M, Gestwa L, Ten Cate WJ, Lautermann J, Brugger H, Maier H, et al. Distinct thyroid hormone-dependent expression of TrkB and p75NGFR in nonneuronal cells during the critical TH-dependent period of the cochlea. *J Neurobiol* 1999;38:338-56.
- Kamiya K, Takahashi K, Kitamura K, Momoi T, Yoshikawa Y. Mitosis and apoptosis in postnatal auditory system of the C3H/He strain. *Brain Res* 2001;901:296-302.
- Raff M. Cell suicide for beginners. *Nature* 1998;396:119-22.
- Miller DK. The role of the Caspase family of cysteine proteases in apoptosis. *Semin Immunol* 1997;9:35-49.
- Nicholson DW, Thornberry NA. Caspases: killer proteases. *Trends Biochem Sci* 1997;22:299-306.
- Liu WJ, Yang J. Preferentially regulated expression of connexin 43 in the developing spiral ganglion neurons and afferent terminals in post-natal rat cochlea. *Eur J Histochem* 2015;59:2464.
- Liu WJ, Yang J. Developmental expression of inositol 1, 4, 5 trisphosphate receptor in the postnatal rat cochlea. *Eur J Histochem* 2015;59:2486.
- Hinojosa R. Note on development of Corti's organ. *Acta Otolaryngol* 1977;84:338-51.
- Tritsch NX, Zhang YX, Ellis-Davies G, Bergles DE. ATP-induced morphological changes in supporting cells of the developing cochlea. *Purinergic Signal* 2010;6:155-66.
- Green DR. Means to an end: apoptosis and other cell death mechanisms. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York; 2011.
- Leon Y, Sanchez-Galiano S, Gorospe I. Programmed cell death in the development of the vertebrate inner ear. *Apoptosis* 2004;9(255-64).
- Nicholson DW. Caspase structure, proteolytic substrates, and function during apoptotic cell death. *Cell Death Differ* 1999;6:1028-42.
- Janicke RU, Ng P, Sprengart ML, Porter AG. Caspase-3 is required for alpha-fodrin cleavage but dispensable for cleavage of other death substrates in apoptosis. *J Biol Chem* 1998;273:15540-5.
- Vaux DL. A boom time for necrobiology. *Cell Biol* 1993;3:877-8.
- Creagh EM. Caspase crosstalk: integration of apoptotic and innate immune signalling pathways. *Trends Immunol* 2014;35:631-40.
- Adrain C, Martin SJ. The mitochondrial apoptosome: a killer unleashed by the cytochrome seas. *Trends Biochem Sci* 2001;26:390-7.
- Vorasubin N, Hosokawa S, Hosokawa K, Ishiyama G, Ishiyama A, Lopez IA. Neuroglobin immunoreactivity in the human cochlea. *Brain Res* 2016;1630:56-63.
- Lu X, Sun S, Qi J, Li W, Liu L, Zhang Y, et al. *Bmi1* regulates the proliferation of cochlear supporting cells via the canonical Wnt signaling pathway. *Mol Neurobiol* 2017;54:1326-39.

Non commercial use only