

DOI: [10.4081/ejh.2018.2890](https://doi.org/10.4081/ejh.2018.2890)

### Distribution of non-myelinating Schwann cells and their associations with leukocytes in mouse spleen revealed by immunofluorescence staining

Bin Ma,<sup>1</sup> Changfu Yin,<sup>2</sup> Dailun Hu,<sup>2</sup> Mark Newman,<sup>1</sup> Philip K. Nicholls,<sup>1</sup> Zhanjun Wu,<sup>3</sup>  
Wayne K. Greene,<sup>1</sup> Zhongli Shi<sup>2</sup>

<sup>1</sup>School of Veterinary and Life Sciences, Murdoch University, Murdoch, WA, Australia


<sup>2</sup>Clinical College, Hebei Medical University, Shijiazhuang, Hebei, China

<sup>3</sup>Institute of Cereal and Oil Crop, Hebei Academy of Agriculture and Forestry Sciences, Shijiazhuang, Hebei, China

**Correspondence:** Zhongli Shi, Clinical College, Hebei Medical University, 309 Jianhuanan Street, Shijiazhuang, Hebei 050031, China. Tel. +86.311.86265948 - Fax: +86.311.86265911. E-mail: [shizhongli@hebmh.edu.cn](mailto:shizhongli@hebmh.edu.cn)

**Key words:** Non-myelinating Schwann cells (NMSC); lymphocyte; dendritic cell (DC); spleen; immunofluorescence staining.

---



**Supplementary Figure 1.** Co-staining of GFAP (green) with other NMSC marker (p75<sup>NTR</sup>, in red) and neuronal markers (PGP9.5 and neurofilament, in red) of spleen sections. The white arrows indicate NMSC/NMSC processes associated with other NMSC or neuronal markers. b) This image is a maximal intensity projection of a Z-stack. CA, central artery; T, trabecula; MZ, marginal zone. Objective lens: 40x; Stack size: 6.0  $\mu\text{m}$ ; optical slice interval: 0.50  $\mu\text{m}$ . Scale bar: 20  $\mu\text{m}$ .