

DOI: [10.4081/ejh.2018.2971](https://doi.org/10.4081/ejh.2018.2971)

### Epidermal growth factor-like domain multiple 7 (EGFL7): Expression and possible effect on biliary epithelium growth in cholangiocarcinoma

Caterina L. Mammola,<sup>1</sup> Antonella Vetuschi,<sup>2</sup> Luigi Pannarale,<sup>1</sup> Roberta Sferra,<sup>2</sup>  
Romina Mancinelli<sup>1</sup>


<sup>1</sup>Department of Anatomical, Histological, Forensic Medicine and Orthopedics Sciences, Sapienza University of Rome

<sup>2</sup>Department of Biotechnological and Applied Clinical Sciences, University of L'Aquila, Italy

**Correspondence:** Romina Mancinelli, Department of Anatomical, Histological, Forensic Medicine and Orthopedics Sciences, Sapienza University of Rome, Via Alfonso Borelli 50, 00161 Rome, Italy. Tel. +39.06.49918063; Fax: +39.06.49918062.

E-mail: [romina.mancinelli@unitoma1.it](mailto:romina.mancinelli@unitoma1.it)

**Key words:** Liver; biliary epithelium; cholangiocytes; cholangiocarcinoma; EGFL7.



### Supplementary Figure 1.

HuH28 intrahepatic CCA cell line highly expresses EGFL7 compared to the other cell lines: the extrahepatic CCA Mz-ChA-1 and the non-malignant H69. Specific immunoreactivity of representative fields is shown in green and nuclei were stained with 4,6-diamidino-2-phenylindole (DAPI; blue). No staining was visible when primary antibodies were replaced with nonimmune serum (negative control on the left). Scale bar: 20  $\mu$ m.